

Escalade^{®2}



Take Down Tough Weeds with Escalade^{®2}

Nothing knocks down unsightly weeds like Escalade^{®2} herbicide. Escalade² combines three powerful active ingredients to control more than 240 broadleaf weeds. It starts working quickly to devastate weeds and help reduce callbacks. Plus, Escalade² can be tank-mixed with most liquid fertilizers and iron.

KEY PERFORMANCE BENEFITS

- Offers the power of three proven herbicides in one convenient pre-mixed formulation
- Delivers quick visual response, devastates weeds and reduces callbacks
- Excellent control of more than 240 broadleaf weeds including dandelion, plantain, oxalis, chickweed and other hard-to-control species
- Unparalleled clover control
- Can be used for treatments in early spring season as well as spot applications
- Compatible with most liquid fertilizers and iron

EPA REG. NO.	228-442
ACTIVE INGR.	2,4-D (39.53%), fluroxypyr (5.9%), dicamba (4.10%)
FORMULATION	liquid
CHEM. FAMILY	phenoxyacetic acid, pyridine and benzoic acid
HRAC NO.	4
MODE OF ACT.	auxin growth regulators
SIGNAL WORD	warning
RESTRICTED	no
PACKAGE SIZE	5 x 1 gal, 2 x 2.5 gal, 30 gal, 250 gal tote



Nufarm

Grow a better tomorrow.

Escalade®2

KEY USES

Ornamental lawns and turf
Roadsides (including aprons and guardrails),
rights-of-way, and other similar non-crop
areas)
Sod farms

KEY WEEDS CONTROLLED

(See product label for complete list)

Bedstraw
Beggartick
Bitterweed
Bitter wintercress
Black-eyed Susan
Buckhorn
Burweed
Buttercup
Carpetweed
Catsear
Chickweed
Chicory
Clover
Cockle
Cocklebur
Dandelion
Dock
Hawkweed
Henbit
Honeysuckle
Horsenettle
Horsetail
Kochia
Knotweed
Lambsquarters
Lespedeza
Mallow
Morningglory
Mustard
Nettle
Oxalis
Pepperweed
Pigweed
Plantain
Primrose
Purslane
Shepherd's-purse
Sowthistle
Spurge
Tansy ragwort
Thistle
Vervain
Vetch
Willow
Woodsorrel
Woolly croton
Yarrow
Yellow rocket

USE SUMMARY

ORNAMENTAL LAWNS AND TURF

Apply 2 to 3 pints of Escalade®2 in 20 to 240 gallons of water per acre (0.75 to 1.1 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet) to control weeds growing in turf planted to bluegrass, fescue, ryegrass, bentgrass (excluding golf course greens and tees), bahia, bermudagrass and zoysia.

On closely mowed golf course fairway bentgrass, apply this product at a maximum rate of 2 pints in 20 to 240 gallons of water per acre (0.75 fluid ounces in 0.5 to 5.5 gallons of water per 1,000 square feet). During conditions which promote turf stress use lower rates, 1 to 1.5 pints in 20 to 240 gallons of water per acre (0.36 to 0.55 fluid ounces in 0.3 to 5.5 gallons per 1,000 square feet). Slight turf yellowing will disappear after one week. To minimize grass injury, a second application should not be made for at least 4 weeks. Do not use on golf course greens or tees.

SOD FARMS

For bluegrass, ryegrass and fescue, apply 2 to 3 pints per acre.

For creeping bentgrass use 1.25 pints per acre.

For common and hybrid bermudagrass, bahia, zoysia and St. Augustinegrass apply 1.5 to 1.8 pints per acre.

Spray volume for sod farm application is 5 to 175 gallons per acre.

NON-TURF AREAS

Mix 2 to 5 pints per acre in adequate water to thoroughly saturate all weeds with spray mixture. This may require a spray volume of 50 to 300 gallons of water per acre. Apply any time between the time when plants come into full leaf (spring) to when the plants begin to go dormant. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 2.25 ounces of this product per gallon of water and spray to thoroughly wet all foliage.

For control of woody plants, apply to both stems and foliage any time from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 4 to 5 pints of this product per acre in adequate water to thoroughly saturate all weeds with the spray mixture. This may require a spray volume of 200 to 600 gallons of water per acre depending upon the height and thickness of the brush.

Mix thoroughly before spraying.

Maximum seasonal application rate to non-crop sites is 4 pounds 2,4-D acid equivalent per acre per application site.

See label for complete directions for use.

